

Comment on “Spin Correlations in the Paramagnetic Phase and Ring Exchange in La_2CuO_4 ”

L. Raymond, G. Albinet

L2MP, 49 rue Joliot Curie BP 146, Université de Provence, 13384 Marseille, Cedex 13, France

A.-M. S. Tremblay

Département de Physique and RQMP, Université de Sherbrooke, Sherbrooke, Québec, J1K 2R1, Canada.

It is shown that the experiments of A.M. Toader, J. P. Goff, M. Roger, N. Shannon, J. R. Stewart, and M. Enderle, Phys. Rev. Lett. **94**, 197202 (2005) do not provide definitive experimental evidence for ring exchange terms in the Hamiltonian of La_2CuO_4 , even though such terms may be present.

In a recent paper, Toader *et al.* [1] claimed to provide definitive experimental evidence for ring exchange terms in the Hamiltonian of La_2CuO_4 by comparing the experimental antiferromagnetic static spin structure factor $S(\mathbf{Q})$ with high-temperature series expansion. Ring-exchange terms arise at intermediate coupling in the effective low-energy theory for the Hubbard model. The ring-exchange parameters deduced from Ref. [1] agree with those found earlier [2], and in terms of the Hubbard model they correspond to $\kappa \equiv t/U = 1/7.5$ with $t = 0.29 \text{ eV}$.

Fig. 1 compares several results with the corresponding data in Fig.3 of Ref. [1]. The high-temperature series expansion of $S(\mathbf{Q})$ in Ref. [1] are in units where $J/t = 2\tilde{J}_2^{(1)}/t = 4\kappa - 64\kappa^3$ and they are represented by the two dashed lines, along with three of the experimental points in the upper right-hand corner. Our Quantum

Monte Carlo (QMC) data for $U = 7.5t$ and $U = 10t$ are given respectively by circles and by squares linked by a dotted line as a guide to the eye. We used the determinantal method with discretization step $\Delta\tau = 1/10$. When the QMC data appear to have reached a size independent value, we give as lower and upper error bars statistical fluctuations on, respectively, the smallest and largest result. The largest system size is typically $L \times L = 12 \times 12$. When the size dependence is still important, the upper error bar is an upper bound obtained by a $1/L$ extrapolation of the values for the two largest system sizes. Except for the lowest temperatures, the results have converged as a function of L . The horizontal scale for QMC is obtained from $J/t = 4t/U \equiv 4\kappa$ with temperature in units of t , so that the scale $J/(2T)$ depends only on the ratio κ . We also show as a dash-dot line the value of $S(\mathbf{Q})$ obtained from high-temperature series expansion of the Heisenberg model, [3] without ring exchange. We take $\hbar = 1$ for both the Heisenberg and QMC results so that the corresponding $S(\mathbf{Q})$ can be compared with QMC on an absolute scale. However, the absolute scale of the results of Ref. [1] is not given, but on a logarithmic scale this is just a shift of the origin that does not change the slope. So $S(\mathbf{Q})$ for the Heisenberg model and for QMC have all been shifted by precisely the same value to compare with Ref. [1]. Clearly, as expected, the larger the value of U in the QMC calculation, the better the agreement with the Heisenberg model. Note however that both values of U are much closer to the Heisenberg result than to the results of Ref. [1], despite the fact that $U = 7.5t$ should correspond to the exchange parameters (including ring) used in that reference.

In Ref. [1] it was argued that agreement with experiment was better with ring exchange than without it because the high-temperature series results dove tail better with the experiment than the results obtained without ring exchange. We have here a counter example since with just a shift of the origin of the vertical axis of either the $U = 7.5t$ or $U = 10t$ results of the Hubbard model we can smoothly join the experimental data. As an added observation, we show in Fig. 1 that when our QMC data (represented by the same symbols as before but this time without error bars) are plotted as a func-




FIG. 1: High temperature series expansion from Ref. [1], lower dashed line is fifth order, higher dashed line is fourth order. Experimental results in upper right-hand corner. Heisenberg model [3] dashed-dotted line. QMC results $U = 7.5t$ circles and $U = 10t$ squares. When no error bars are indicated, there is additional scaling, as discussed in the text.

tion of $J/t = 2\tilde{J}_2^{(1)}/t = 4\kappa - 64\kappa^3$, as suggested in Ref. [1], then they fall close to the Heisenberg curve and they all join smoothly the experiment.

The experimental results on $S(\mathbf{Q})$ are in the universal regime that can be described by the non-linear sigma model and hence they are insensitive to microscopic details, as noted in Ref. [1]. Our results suggest that unless this $S(\mathbf{Q})$ can be measured in the non-universal regime at higher temperature, it cannot lead to an accurate value of U/t (and hence of ring exchange contribution) by a smoothness argument, especially if additional parameters such as next-nearest-neighbor hopping are taken into account. Experiments involving measurement of $S(\mathbf{Q},\omega)$ at higher energies and shorter wavelengths are more informative, as discussed in Ref. [2], but even in that case it is important to also take into account detailed infor-

mation on the band structure, including further neighbor hoppings, to obtain reliable values of U and consequent ring exchange terms.[4]

We are indebted to M. Gingras, J.-Y. Delannoy, M. Roger and N. Shannon for useful conversations.

-
- [1] A.M. Toader, J. P. Goff, M. Roger, N. Shannon, J. R. Stewart, and M. Enderle, Phys. Rev. Lett. **94**, 197202 (2005).
 - [2] R. Coldea, *et al.* Phys. Rev. Lett. **86**, 5377 (2001).
 - [3] N. Elstner, R.R.P. Singh, and A.P. Young, J. Appl. Phys. **75**, 5943 (1994).
 - [4] J.Y. Delannoy, PhD thesis, ENS Lyon, (unpublished).